

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 21 September 1998	3. REPORT TYPE AND DATES COVERED Final: 15 Feb 96 - 31 Dec 97	
4. TITLE AND SUBTITLE  Modeling the wind and buoyancy driven circulation and ice interaction in the Okhotsk Sea			5. FUNDING NUMBERS  (G)rant N00014-96-1-0463	
6. AUTHOR(S)  Jiayan Yang				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Woods Hole Oceanographic Institution Mail Stop #21 Woods Hole, Massachusetts 02543			8. PERFORMING ORGANIZATION REPORT NUMBER  14046300	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)  Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217-5660  Attn: Michael Van Woert ONR Code 322HL			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES  Attn: Boston Regional Office Admin. Grants Officer				
12a. DISTRIBUTION / AVAILABILITY STATEMENT  Approved for Public Release, Distribution is Unlimited			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  The main goal of this objective is to use coupled ocean-ice numerical models to study the basin scale ocean circulation and ocean-ice interaction in the Okhotsk Sea, and to examine important processes that maintain the unique oceanographic setting, such as the ODTL in this large marginal sea. We use a hierarchy of models, from a basin-averaged column model of ocean-ice interactions (Yang and Honjo, 1996) to a 3-D primitive equation ocean general circulation models coupled with a dynamical and thermodynamical sea-ice model. Meanwhile, we have analyzed some unpublished in situ observations (mainly hydrography records from the Japanese Hydrography Office), satellite SSM/I data of sea-ice concentration, and meteorological observations (from Comprehensive Ocean and Atmosphere Data Sets (COADS) and from the Japan Meteorology Agency) to examine ocean-ice interactions and mechanisms for interannual variations.				
14. SUBJECT TERMS  Okhotsk Dichothermal Layer (ODTL); Sea Ice; Ocean Circulation.			15. NUMBER OF PAGES  6	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT  UL	

## Final Report

Grant No. N00014-96-1-0463

**Title:** Modeling the wind and buoyancy driven circulation and ice interaction in the Okhotsk Sea

**Principal Investigator:** Jiayan Yang

## Abstract

The Sea of Okhotsk is one of the largest semi-enclosed marginal sea, connected with the North Pacific Ocean through numerous Kuril straits, and with the Sea of Japan (the East Sea) through Soya Strait. Despite its moderate geographic location, the Sea of Okhotsk is mostly covered by sea ice in winter. The sea surface temperature in summer, however, reaches about 15°C. This strong seasonal variation gives rise to some important physical oceanography features, such as the Okhotsk Sea Dichothermal Layer (ODTL) (a near-freezing subsurface layer that exists through the whole year) and the formation of the North Pacific Intermediate Water. Meanwhile, satellite observation shows that interannual variation of sea-ice extent is among the greatest in the world. The main objective is to use coupled ocean-ice numerical models to study the basin scale ocean circulation and ocean-ice interaction in the Okhotsk Sea, and to examine important processes that maintain the unique oceanographic setting, such as the ODTL in this large marginal sea. We use a hierarchy of models, from a basin-averaged column model of ocean-ice interactions (Yang and Honjo, 1996) to a 3-D primitive equation ocean general circulation models coupled with a dynamical and thermodynamical sea-ice model. Meanwhile, we have analyzed some unpublished in situ observations (mainly hydrography records from the Japanese Hydrography Office), satellite SSM/I data of sea-ice concentration, and meteorological observations (from Comprehensive Ocean and Atmosphere Data Sets (COADS) and from the Japan Meteorology Agency) to examine ocean-ice interactions and mechanisms for interannual variations.

19980928 039

## Summary





We have developed a basin-averaged coupled ocean-ice model to study the formation and variation of the ODTL and sea-ice cover. The results show that the water Okhotsk Dichothermal Layer is renewed by winter convective mixing when brine rejection associated with sea-ice formation destabilizes the water column. In the summer, the low salinity (due to spring melting of sea ice) and high temperature water in the mixed layer prevents the vertical mixing to reach the dichothermal layer, and therefore, the near freezing layer is able to sustain. Our study also shows that the seemingly net-zero-gain process of winter freezing and summer melting can result in interannual variations in the ODTL and in sea-ice extent. Our study indicates that the strong seasonal variation in the atmosphere forcing fields is primarily responsible to some of the unique observed features in the Okhotsk Sea, such as the existence of the near freezing ODTL. We used this model to study the effects of inflows from Sea of Japan through Soya Strait and from the North Pacific Ocean through Kurile Straits on the mean hydrographic structure and ODTL variability. The role of fresh water flux from Amur River runoff and surface evaporation-precipitation is also investigated. The result was published in *Journal of Geophysical Research* in 1996.

We have extended our study to using a 3-D model so that key processes can be examined in a more realistical setting. We use this model to study the mean circulation and spatial variations of sea-ice extent. We are focused on modeling the seasonal cycle but will extend to include interannual variations. Dr. S. Honjo and I have been working on some newly available hydrography and geochemical tracer data (unpublished previously) collected by R/V Shoyo (Shoyo cruise 9007 and 9105) of the Hydrography Office of Japan during August, 1990 and 1991. Significant changes in potential temperature, salinity and dissolved oxygen content were observed between 1990 and 1991. We therefore analyzed the satellite sea-ice observations and COADS meteorological data to examine the causes for these interannual variations. A manuscript, co-authored with S. Honjo and entitled "Atmosphere-Ocean-Ice Interactions in the Okhotsk Sea", has been written based on these analyses.

Meanwhile, we have examined observations of hydrography, tracers, surface wind and air temperature, and satellite sea-ice concentration to study the responses of a coupled ocean-ice system to variations in the atmosphere forcings. It was found that, the ODTL water in the central Okhotsk basin, between 50N and 55.4N in latitude, was considerably warmer and fresher in 1991 than in 1990. Figure 1 shows the potential temperature in these two sections in 1991 and 1990. The ODTL structure is clearly shown in all sections. The difference between 1990 and 1991 are considerable in both surface layer and in the ODTL (Figure 2). The COADS data showed that the northwesterly wind from Siberia was much stronger in the winter of 1990 than in 1991. Therefore, the air temperature was more than 5 deg C colder over most of the basin in January of 1990 than in the same month in 1991. The satellite SSM/I data showed that the sea-ice concentration in the central basin was up to 60(Figure 3). Therefore, the more vigorous convective mixing in 1990 was likely the cause for the observed difference in potential temperature and salinity. The Yang and Honjo (1996) model confirmed our analyses. This indicates that the coupled ocean-ice system in the Okhotsk Sea is very sensitive to variations in the atmosphere, especially in the winter season when convective mixing occurs. It also suggests that a predictive skill can be developed that can be used to infer the oceanographic conditions based on winter atmospheric condition.

## Reference

- Yang, J. and S. Honjo, 1996: Modeling the near-freezing dichothermal layer in the Sea of Okhotsk and its interannual variations. *Journal of Geophysical Research*, **101**, 16,421-16,433.
- Yang, J. and S. Honjo. Atmosphere-Ocean-Ice Interactions in the Okhotsk Sea. Manuscript.






Figure 2(a) Temperature Difference (1991-1990) at 149°15'W






Figure 2(b) Temperature Difference (1991-1990) at 149°50'W

Sea-Ice Concentration in the Okhotsk Sea (March, 1990)



Sea-Ice Concentration in the Okhotsk Sea (March, 1991)



Sea-Ice Concentration Difference: 1990-1991 (March)

